

Exceptional values in holomorphic families of entire functions

Alexandre Eremenko*

March 27, 2005

In 1926, Julia [5] studied singularities of implicit functions defined by equations $f(z, w) = 0$, where f is an entire function of two variables. Among other things, he investigated the exceptional set P consisting of those w for which such equation has no solutions z . In other words, P is the complement of the projection of the analytic set $\{(z, w) : f(z, w) = 0\}$ onto the second coordinate. Julia proved that P is closed and cannot contain a continuum, unless it coincides with w -plane. Lelong [6] and Tsuji [12], [13, Thm. VIII.37] independently improved this result by showing that the logarithmic capacity of P is zero if $P \neq \mathbf{C}$. In the opposite direction, Julia [5] proved that every discrete set $P \subset \mathbf{C}$ can occur as the exceptional set. He writes: “*Resterait à voir si cet ensemble, sans être continu, peut avoir la puissance du continu*”.¹

According to Alan Sokal (private communication), the same question arises in holomorphic dynamics, when one tries to extend to holomorphic families of transcendental entire functions the results of Lyubich [8] on holomorphic families of rational functions.

In this paper, we show that in general, the result of Lelong and Tsuji is best possible: every closed set of zero capacity can occur as an exceptional set (Theorem 1). Then we study a related problem of dependence of Picard exceptional values of the function $z \mapsto f(z, w)$ on the parameter w (Theorem 2).

It is known that the exceptional set P is discrete in the important case that $z \mapsto f(z, w)$ are functions of finite order. This was discovered by Lelong

*Supported by NSF grants DMS-0100512 and DMS-0244547

¹It remains to see whether this set, without being a continuum, can have the power of continuum.

in [6] and later the result was generalized to the case of multi-dimensional parameter w in [7, Thm. 3.44].

We also mention that the set P has to be analytic in certain holomorphic families of entire functions with finitely many singular values, considered in [1, 2]. These families may consist of functions of infinite order.

We begin with a simple proof of a version of Lelong's theorem on functions of finite order.

Proposition 1. *Let D be a complex manifold, and $f : \mathbf{C} \times D \rightarrow \mathbf{C}$ an analytic function, such that the entire functions $z \mapsto f(z, w)$ are not identically equal to zero and are of finite order for all $w \in D$. Then the set*

$$P = \{w \in D : (\forall z \in \mathbf{C}) f(z, w) \neq 0\} \quad (1)$$

is analytic.

Corollary. *Let D be a region in $\mathbf{C}$, and $f : \mathbf{C} \times D \rightarrow \mathbf{C}$ an analytic function, such that entire functions $z \mapsto f(z, w)$ are of finite order for all $w \in D$. Then the set P as in (1) is discrete or $P = D$.*

Indeed, the set $A = \{w \in D : f(., w) \equiv 0\}$ is discrete (unless $f = 0$ when there is nothing to prove). So there exists a function g holomorphic in D whose zero set is A , and such that f/g satisfies all conditions of Proposition 1. $\square$

Remarks.

1. In general, when D is of dimension greater than one, and the set A is not empty, one can only prove that $A \cup P$ is contained in a proper analytic subset of D , unless $A \cup P = D$, [7, Thm 3.44].

2. If the order of $f(., w)$ is finite for all $w \in D$ then this order is bounded on compact subsets of D [7, Thm. 1.41].

Proof of Proposition 1. We assume without loss of generality that

$$f(0, w) \neq 0 \quad \text{for } w \in D. \quad (2)$$

(shift the origin in $\mathbf{C}$ and shrink D , if necessary), and that the order of the function $f(., w)$ does not exceed λ for all $w \in D$ (see Remark 2 above).

Let p be an integer, $p > \lambda$. Then, for each w , f has the Weierstrass representation

$$f(z, w) = e^{c_0 + \dots + c_p z^p} \prod_{a: f(a, w)=0} \left(1 - \frac{z}{a}\right) e^{z/a + \dots + z^p/pa^p},$$

where a are the zeros of $f(., w)$ repeated according to their multiplicities, and c_j and a depend on w . Taking the logarithmic derivative, differentiating it p times, and substituting $z = 0$, we obtain for each $w \in D$:

$$F_p(w) = \frac{d^p}{dz^p} \left(\frac{df}{f dz} \right) \Big|_{z=0} = p! \sum_{a: f(a,w)=0} a^{-p-1}.$$

The series in the right hand side is absolutely convergent because of our choice of p . The functions F_p are holomorphic in D , in view of (2). Clearly $w \in P$ implies $F_p(w) = 0$ for all $p > \lambda$. In the opposite direction, $F_p(w) = 0$ for all $p > \lambda$ means that all but finitely many derivatives with respect to z at $z = 0$ of the function $df/f dz$ meromorphic in $\mathbf{C}$ are equal to zero, so this meromorphic function is a polynomial, and thus $f(z, w) = \exp(c_0 + \dots + c_p z^p)$, that is $w \in P$. $\square$

The following result is due to Lelong and Tsuji but they state it only for the case $\dim D = 1$, and we need a multi-dimensional version in the proof of Theorem 2 below.

Proposition 2. *Let D be a connected complex manifold, and $f : \mathbf{C} \times D \rightarrow \mathbf{C}$ an analytic function, such that the entire functions $z \mapsto f(z, w)$ are not identically equal to zero. Then the set*

$$P = \{w \in D : (\forall z \in \mathbf{C}) f(z, w) \neq 0\}$$

is closed and pluripolar or coincides with D .

Proof. Suppose that $P \neq D$. It is enough to show that every point $w_0 \in D$ has a neighborhood U such that $P \cap U$ is pluripolar. By shifting the origin in $\mathbf{C}_z$ and shrinking U we may assume that $f(0, w) \neq 0$ for all $w \in U$. Let $r(w)$ be the smallest of the moduli of zeros of the entire function $z \mapsto f(z, w)$. If this function has no zeros, we set $r(w) = +\infty$. We are going to prove that $\log r$ is a continuous plurisuperharmonic function.

First we prove continuity of $r : U \rightarrow (0, +\infty]$. Indeed, suppose that $r(w_0) < \infty$, and let k be the number of zeros of $f(., w_0)$ on $|z| = r(w_0)$, counting multiplicity. Let $\epsilon > 0$ be so small that the number of zeros of $f(., w_0)$ in $|z| \leq r(w_0) + \epsilon$ equals k . Then

$$\int_{|z|=r(w_0) \pm \epsilon} \frac{df}{f dz} dz = \begin{cases} 2\pi i k, \\ 0. \end{cases}$$

As the integrals depend on w continuously, we conclude that r is continuous at w_0 . Consideration of the case $r(w_0) = +\infty$ is similar.

Now we verify that the restriction of $\log r$ to any complex line is superharmonic. Let $\zeta \rightarrow w(\zeta)$ be the equation of such line, where $w(0) = w_0$. Let z_0 be a zero of $f(\cdot, w_0)$ of the smallest modulus. We verify the inequality for the averages of $\log r$ over the circles $|\zeta| = \delta$, where δ is small enough. According to the Weierstrass Preparation theorem, the set $Q = \{(z, \zeta) : f(z, w(\zeta)) = 0\}$ is given in a neighborhood of $(z_0, 0)$ by an equation of the form

$$(z - z_0)^p + b_{p-1}(\zeta)(z - z_0)^{p-1} + \dots + b_0(\zeta) = 0,$$

where b_j are analytic functions in a neighborhood of 0, $b_j(0) = 0$. We rewrite this as

$$z^p + c_{p-1}(\zeta)z^{p-1} + \dots + c_0(\zeta) = 0, \quad (3)$$

where c_j are analytic functions, and

$$c_0(0) = (-z_0)^p. \quad (4)$$

Let V be a punctured disc around 0; we choose its radius so small that $c_0(\zeta) \neq 0$ in V . For ζ in V , let $z_i(\zeta)$, $i = 1, \dots, p$ be the branches of the multi-valued function $z(\zeta)$ defined by equation (3). Then

$$\begin{aligned} \frac{1}{2\pi} \int_{-\pi}^{\pi} \log r(w(\delta e^{i\theta})) d\theta &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \min_i \log |z_i(\delta e^{i\theta})| d\theta \\ &\leq \frac{1}{2\pi p} \int_{-\pi}^{\pi} \log \prod_i |z_i(\delta e^{i\theta})| d\theta = \frac{1}{2\pi p} \int_{-\pi}^{\pi} \log |c_0(\delta e^{i\theta})| d\theta \\ &= \log |z_0| = \log r(w_0), \end{aligned}$$

where we used (4) and harmonicity of $\log |c_0|$ in $V \cup \{0\}$. This completes the proof of plurisuperharmonicity. As $P = \{w : \log r(w) = +\infty\}$ we conclude that P is pluripolar. $\square$

Our first theorem answers the question of Julia; it shows that the restriction of finiteness of order cannot be removed in Proposition 1, and that Proposition 2 is best possible, at least when $\dim D = 1$.

Theorem 1. *Let $D \subset \mathbf{C}$ be the unit disc, and P an arbitrary compact subset of D of zero capacity. Then there exists a holomorphic function $f : \mathbf{C} \times D \rightarrow$*

$\mathbf{C}$, such that for every $w \in P$ the equation $f(z, w) = 0$ has no solutions, and for each $w \in D \setminus P$ it has infinitely many solutions.

It is not clear whether a similar result holds with multidimensional parameter space D and arbitrary closed pluripolar set $P \subset D$.

Proof. Let $\phi : D \rightarrow D \setminus P$ be a universal covering. Let S be the set of singular points of ϕ on the unit circle. Then S is a closed set of zero Lebesgue measure.

(We recall a simple proof of this fact. As a bounded analytic function, ϕ has radial limits almost everywhere. It is easy to see that a point where the radial limit has absolute value 1 is not a singular point of ϕ . Thus the radial limits exist and belong to P almost everywhere on S . Let u be the “Evans potential” of P , that is a harmonic function in $D \setminus P$, continuous in $\overline{D}$ and such that $u(\zeta) = 0$ for $\zeta \in \partial D$ and $u(\zeta) = -\infty$ for $\zeta \in P$. Such function exists for every compact set P of zero capacity. Now $v = u \circ \phi$ is a negative harmonic function in the unit disc, whose radial limits on S are equal to $-\infty$, thus $|S| = 0$ by the classical uniqueness theorem.)

According to a theorem of Fatou, (see, for example, [3, Ch. VI]), for every set S of zero Lebesgue measure on ∂D , there exists a holomorphic function g in D , continuous in $\overline{D}$ and such that

$$\{\zeta \in \overline{D} : g(\zeta) = 0\} = S.$$

In particular, g has no zeros in D .

Now we define the following set $Q \subset \mathbf{C} \times D$:

$$Q = \{(1/g(\zeta), \phi(\zeta)) : \zeta \in D\}.$$

It is evident that the projection of Q on the second coordinate equals $D \setminus P$. It remains to prove that the set Q is analytic. For this, it is enough to establish that the map

$$\Phi : D \rightarrow \mathbf{C} \times D, \quad \Phi(\zeta) = (1/g(\zeta), \phi(\zeta))$$

is proper. Let $K \subset \mathbf{C} \times D$ be a compact subset. Then the closure of $\Phi^{-1}(K)$ in $\overline{D}$ is disjoint from S because g is continuous and $1/g(\zeta) \rightarrow \infty$ as $\zeta \rightarrow S$. On the other hand, for every point $\zeta \in \partial D \setminus S$, the limit

$$\lim_{\zeta \rightarrow \zeta_0} \phi(\zeta)$$

exists and has absolute value 1. So $\Phi^{-1}(K)$ is compact in D .

Now the existence of the required function f follows from the solvability of the Second Cousin problem [4, sect. 5.6]. $\square$

Notice that the map Φ constructed in the proof is an immersion.

We recall that a point $a \in \mathbf{C}$ is called an exceptional value of an entire function f if the equation $f(z) = a$ has no solutions. Picard's Little theorem says that a non-constant entire function can have at most one exceptional value.

Let f be an entire function of z depending of the parameter w holomorphically, as in Propositions 1 and 2, and

assume in the rest of the paper that for all $w \in D$, $f(., w) \neq \text{const}$.

Let $n(w) \in \{0, 1\}$ be the number of exceptional values of $f(., w)$.

Question 1. What can be said about $n(w)$ as a function of w ?

Example 1. $f(z, w) = e^z + wz$. We have $n(0) = 1$ and $n(w) = 0$ for $w \neq 0$.

Example 2. $f(z, w) = (e^{wz} - 1)/w$ for $w \neq 0$, and $f(z, 0) = z$. We have $n(0) = 0$, while $n(w) = 1$ for $w \neq 0$. The exceptional value $a(w) = -1/w$ tends to infinity as $w \rightarrow 0$.

Thus n is neither upper nor lower semicontinuous.

Example 3. Let

$$f(z, w) = \int_{-\infty}^z (\zeta + w)e^{-\zeta^2/2} d\zeta,$$

where the contour of integration consists of the negative ray, passed left to right, followed by a curve from 0 to z . We have $f(z, 0) = -e^{-z^2/2}$, which has exceptional value 0, so $n(0) = 1$. It is easy to see that there are no exceptional values for $w \neq 0$, so $n(w) = 0$ for $w \neq 0$. Thus $n(w)$ is the same as in Example 1, but this time we have an additional feature that the set of singular values of $f(., w)$ is finite for all $w \in \mathbf{C}$, namely, there is one critical value $f(-w, w)$ and two asymptotic values, 0 and $\sqrt{2\pi}w$.

Question 2. Suppose that $n(w) \equiv 1$, and let $a(w)$ be the exceptional value of $f(., w)$. What can be said about $a(w)$ as a function of w ?

For functions of finite order, Question 2 was addressed by Nishino [9] who proved the following:

Let f be an entire function of two variables such that $z \mapsto f(z, w)$ is a non-constant function of finite order for all w . If $n(w) = 1$ for all w in some set having a finite accumulation point, then there exists a meromorphic function $\tilde{a}(w)$ such that $a(w) = \tilde{a}(w)$ when $a(w) \neq \infty$, and $f(\cdot, w)$ is a polynomial when $\tilde{a}(w) = \infty$.

Example 4. (Nishino) Let $f(z, w) = (e^{we^z} - 1)/w$, $w \neq 0$ and $f(z, 0) = e^z$. Then f is an entire function of two variables and $n(w) \equiv 1$. However $a(w) = -1/w$, $w \neq 0$ and $a(0) = 0$, so a is a discontinuous function of w .

Nishino also proved that for arbitrary entire function f of two variables, with $n(w) = 1$ in some region D , the set of discontinuity of the function $a(w)$ is closed and nowhere dense in D .

Our Theorem 2 below gives a complete answer to Question 2. We first prove the following semi-continuity property of the set of exceptional values which holds for all *meromorphic functions* holomorphically depending on parameter. We always assume that $z \mapsto f(z, w)$ is non-constant for all w . Denote

$$A(w) = \{a \in \overline{\mathbf{C}} : (\forall z \in \mathbf{C}) f(z, w) \neq a\}.$$

Proposition 3. *For every $w_0 \in D$ and every $\epsilon > 0$ there exists $\delta > 0$ such that $|w - w_0| < \delta$ implies that $A(w)$ is contained in the ϵ -neighborhood of $A(w_0)$ with respect to the spherical metric.*

Proof. Let U be the open ϵ -neighborhood of $A(w_0)$. Then $K = \overline{\mathbf{C}} \setminus U$ is compact, so there exists $r > 0$ such that the image of the disc $|z| < r$ under $f(\cdot, w_0)$ contains K . Then by Hurwitz's theorem, for every w close enough to w_0 the image of the disc $|z| < r$ under $f(\cdot, w)$ will also contain K . $\square$

Corollary 1. *The set of meromorphic functions having no exceptional values on the Riemann sphere is open in the topology of uniform convergence on compact subsets of $\mathbf{C}$ with respect to the spherical metric in the image.*

The set of entire functions whose only exceptional value is ∞ is not open as Example 2 above shows.

Corollary 2. *Suppose that $f(\cdot, w)$ is entire and has an exceptional value $a(w) \in \mathbf{C}$ for all w on some subset $E \subset D$. If $a(w)$ is bounded on E then its restriction on E is continuous.*

Example 4 shows that a can be discontinuous.

Theorem 2. *Let $f : \mathbf{C} \times D \rightarrow \mathbf{C}$, be a holomorphic function, where D is a region in $\mathbf{C}$, and $z \mapsto f(z, w)$ is not constant for all $w \in D$. Assume that for some function $a : D \rightarrow \mathbf{C}$ we have $f(z, w) \neq a(w)$ for all $z \in \mathbf{C}$.*

Then there exists a discrete set $E \subset D$ such that a is holomorphic in $D \setminus E$, and $a(w) \rightarrow \infty$ as $w \rightarrow w_0$, for every $w_0 \in E$.

So the singularities of a can be only of the type described in Example 4.

The main ingredient of the proof of Theorem 2 is the following recent result of N. Shcherbina [11]:

Theorem A. *Let h be a continuous function in a region $G \in \mathbf{C}^n$. If the graph of h is a pluripolar subset of $\mathbf{C}^{n+1}$ then h is analytic.*

Proof of Theorem 2. Consider the analytic set

$$Q = \{(z, w, a) \in \mathbf{C}_z \times D \times \mathbf{C}_a : f(z, w) - a = 0\}.$$

Let R be the complement of the projection of Q onto $D \times \mathbf{C}_a$. Then it follows from the assumptions of Theorem 2 and Picard's theorem that R is the graph of the function $w \mapsto a(w)$. So R is a non-empty proper subset of $D \times \mathbf{C}_a$. Proposition 1 implies that R is closed in D .

According to Proposition 2, R is pluripolar.

It follows from Proposition 3 that $w \mapsto |a(w)|$ is lower semi-continuous, so the sets

$$E_n = \{w \in D : |a(w)| \leq n\}, \quad \text{where } n = 1, 2, 3, \dots$$

are closed. We have $E_1 \subset E_2 \subset \dots$, and the assumptions of Theorem 2 imply that $D = \cup E_j$. Let E_j^0 be the interiors of E_j , then $E_1^0 \subset E_2^0 \subset \dots$, and the set $G = \cup E_j^0$ is open.

We claim that G is dense in D . Indeed, otherwise there would exist a disc $U \subset D$ which is disjoint from G . But then the closed sets $E_j \setminus E_j^0$ with empty interiors would cover D , which is impossible by the Baire category theorem. This proves the claim.

As a is locally bounded on G , Corollary 2 of Proposition 3 implies that a is continuous in G , so by Shcherbina's theorem, a is analytic in G . It follows from the definition of G that a does not have an analytic continuation from any component of G to any boundary point of G . Our goal is to prove that $D \setminus G$ consists of isolated points.

If G has an isolated boundary point w_0 then we have

$$\lim_{w \rightarrow w_0} a(w) = \infty. \quad (5)$$

Indeed, by Proposition 3, the limit set of $a(w)$ as $w \rightarrow w_0$, $w \in G$ consists of at most two points, $a(w_0)$ and ∞ . On the other hand, this limit set is connected. So the limit exists. If the limit is finite, then it is equal to $a(w_0)$ and the Removable Singularity Theorem gives an analytic continuation of a to $G \cup w_0$, contradicting the statement above that there is no such continuation. So the limit is infinite and (5) holds.

We add to G all its isolated boundary points, thus obtaining new open set G' containing G . Our function a has a *meromorphic* continuation to G' , which we call $\tilde{a}$. This meromorphic continuation coincides with a in G but does not coincide at the added points $G' \setminus G$.

We claim that G' has no isolated boundary points. Indeed, suppose that w_0 is an isolated boundary point of G' . By the same argument as above, the limit (5) exists and is infinite. Then $\tilde{a}$ can be extended to w_0 such that the extended function has a pole at w_0 , but then w_0 would be an isolated boundary point of G (poles cannot accumulate to a pole) so $w_0 \in G'$ by definition of G' . This contradiction proves the claim.

Let F' be the complement of G' with respect to D . Then F' is closed, nowhere dense subset of D . Furthermore, F' has no isolated points, because such points would be isolated boundary points of G' . So F' is perfect or empty. Our goal is to prove that F' is empty.

Assume the contrary, that is that F' is perfect. The closed sets E_n cover the locally compact space F' , so by the Baire category theorem one of these E_n contains a relatively open part of F' . This means that there exists a positive integer n and an open disc $U \subset D$ intersecting F' and such that

$$|a(w)| \leq n \quad \text{for } w \in U \cap F'. \quad (6)$$

By Corollary 2 of Proposition 3, this implies that the restriction of a on $U \cap F'$ is continuous.

We are going to prove that $\tilde{a}$ has a continuous extension from G' to $U \cap F'$, and this extension agrees with the restriction of a on $U \cap F'$. Let W be a point of $U \cap F'$, and (w_k) a sequence in G' tending to W . Choosing a subsequence we may assume that there exists a limit

$$\lim_{k \rightarrow \infty} \tilde{a}(w_k), \quad (7)$$

finite or infinite. By a small perturbation of the sequence that does not change the limit of $\tilde{a}(w_k)$ we may assume that w_k are not poles of $\tilde{a}$ so $a(w_k) = \tilde{a}(w_k)$. By Proposition 3, the limit (7) can only be $a(W)$ or ∞ .

To prove continuity we have to exclude the latter case. So suppose that

$$\lim_{k \rightarrow \infty} \tilde{a}(w_k) = \infty. \quad (8)$$

Let C_k be the component of the set

$$G' \cap \{w \in U : |w - W| < 2|w_k - W|\},$$

that contains w_k , and let

$$m_k = \inf\{|\tilde{a}(w)| : w \in C_k\}. \quad (9)$$

We claim that $m_k \rightarrow \infty$. Indeed, suppose this is not so, then choosing a subsequence we may assume that $m_k \leq m$ for some $m > n$. Then there exists a curve in C_k connecting w_k to some point $w'_k \in C_k$ such that $|\tilde{a}(w'_k)| \leq m+1$. As $|a|$ is continuous in C_k , (8) implies that this curve contains a point y_k such that $|\tilde{a}(y_k)| = m+1$. By selecting another subsequence we achieve that

$$\lim_{k \rightarrow \infty} \tilde{a}(y_k) = y, \quad \text{where } |y| = m+1 > n.$$

As $|a(W)| \leq n$, we obtain a contradiction with Proposition 3. This proves our claim that $m_k \rightarrow \infty$ in (9).

So we can assume that

$$m_k \geq n+1 \quad \text{for all } k. \quad (10)$$

Let us show that this leads to a contradiction. Fix k , and consider the limit set of $\tilde{a}(w)$ as $w \rightarrow \partial C_k \cap U$ from $C_k \cap U$. In view of (6), (10) and Proposition 3, this limit set consists of the single point, namely ∞ . To see that this is impossible, we use the following

Lemma. *Let V and C be two intersecting regions in $\mathbf{C}$, and g a meromorphic function in C such that*

$$\lim_{w \rightarrow W, w \in C} g(w) = \infty \quad \text{for all } W \in V \cap \partial C.$$

Then $V \cap \partial C$ consists of isolated points in V and g has meromorphic extension from C to $V \cup C$.

Proof. By shrinking V , we may assume that $|g(w)| \geq 1$ for $w \in C \cap V$. Then $h = 1/g$ has a continuous extension from C to $C \cup V$ by setting $h(w) = 0$ for $w \in V \setminus C$. The extended function is holomorphic on the set

$$\{w \in C \cap V : h(w) \neq 0\},$$

so by Rado's theorem [10, Thm. 3.6.5], h is analytic in $V \cup C$, $1/g$ gives the required meromorphic extension of g . $\square$

Applying this Lemma with

$$C = C_k, V = \{w \in U : |w - W| < 2|w_k - W|\}$$

and $g = \tilde{a}$, and taking into account that $\partial C_k \cap V \subset F'$, and F' has no isolated points, we arrive at a contradiction which completes the proof that $\tilde{a}$ has a continuous extension to $F' \cap U$, an extension which agrees with a on $F' \cap U$.

From Theorem A we obtain now that a is analytic on $F' \cap U$, which contradicts the fact stated in the beginning of the proof that a has no analytic continuation from G .

This contradiction shows that $F' = \emptyset$ which proves the theorem. $\square$

The author thanks Alan Sokal for asking the question which is the subject of this paper, and Adam Epstein, Laszlo Lempert, Pietro Poggi-Corradini, Alexander Rashkovskii and Nikolay Scherbina for stimulating discussions.

References

- [1] A. E. Eremenko and M. Yu. Lyubich, Structural stability in some families of entire functions, *Funct. Anal. Appl.* 19 (1985) 323–324.
- [2] A. E. Eremenko and M. Yu. Lyubich, Dynamical properties of some classes of entire functions, *Ann. Inst. Fourier, Grenoble* 42 (1992) 989–1020.
- [3] Kenneth Hoffman, *Banach spaces of analytic functions*, Prentice-Hall, Englewood Cliffs, N. J. 1962.
- [4] L. Hörmander, *An introduction to complex analysis in several variables*, D. van Nostrand, Princeton, NJ, 1966.

- [5] Gaston Julia, Sur le domaine d'existence d'une fonction implicite définie par une relation entière $G(x, y) = 0$, Bull. soc. math. France 54 (1926) 26-37; CRAS 182 (1926) 556.
- [6] P. Lelong, Sur les valeurs lacunaires d'une relation a deux variables, Bull. Sci. math. 2^e série, t. 66 (1942) 103–108, 112–125.
- [7] P. Lelong and L. Gruman, Entire functions of several complex variables, Springer, Berlin, 1986.
- [8] M. Yu. Lyubich, Investigation of the stability of the dynamics of rational functions, Teor. Funktsii Funktsional. Anal. i Prilozhen. No. 42 (1984) 72–91. (Russian.) Translated in Selecta Math. Soviet. 9 (1990) no. 1 69–90.
- [9] Toshio Nishino, Sur les valeurs exceptionnelles au sens de Picard d'une fonction entière de deux variables, J. Math. Kyoto Univ., 2-3 (1963) 365–372.
- [10] Thomas Ransford, Potential theorem in the complex plane, Cambridge UP, Cambridge 1995.
- [11] N. V. Shcherbina, Pluripolar graphs are holomorphic,
<http://xxx.lanl.gov/abs/math.CV/0301181>, accepted in Acta math.
- [12] Masatsugu Tsuji, On the domain of existence of an implicit function defined by an integral relation $G(x, y) = 0$, Proc. Imperial Academy 19 (1943) 235–240.
- [13] M. Tsuji, Potential theory in modern function theory, Maruzen, Tokyo 1959.

Purdue University, West Lafayette, IN, U.S.A.
eremenko@math.purdue.edu